arXiv:1511.06782v1 [math.CO] 20 Nov 2015

Pseudoachromatic and connected-pseudoachromatic indices of the complete graph *

M. Gabriela Araujo-Pardo
garaujo@matem.unam.mx

Christian Rubio-Montiel
christian@matem.unam.mx

Instituto de Matemáticas
Universidad Nacional Autónoma de México
Ciudad Universitaria, 04510, México D.F.

November 24, 2015

Abstract

A *complete k-coloring* of a graph G is a (not necessarily proper) k-coloring of the vertices of G such that each pair of different colors appears in an edge. A complete k-coloring is also called *connected*, if each color class induces a connected subgraph of G. The *pseudoachromatic index* of a graph G, denoted by $\psi'(G)$, is the largest k for which the line graph of G has a complete k-coloring. Analogously the *connected-pseudoachromatic index* of G, denoted by $\psi'_c(G)$, is the largest k for which the line graph of G has a connected and complete k-coloring.

In this paper, we study these two parameters for the complete graph K_n. Our main contribution is to improve the linear lower bound for the connected pseudoachromatic index given by Abrams and Berman [Australas J Combin 60 (2014), 314–324] and provide an upper bound. These two bounds prove that, for any integer $n \geq 8$, the order of $\psi'_c(K_n)$ is $n^{3/2}$.

Related to the pseudoachromatic index, we prove that, for q a power of 2 and $n = q^2+q+1$, $\psi'(K_n)$ is at least $q^3 + 2q - 3$ which improves the bound $q^3 + q$ given by Araujo, Montellano and Strausz [J Graph Theory 66 (2011), 89–97].

1 Introduction

Let $G = (V(G), E(G))$ a finite simple graph. A *complete* k-coloring of G is an assignment $\varsigma\colon V(G) \to [k]$ (where $[k] := \{1, \ldots, k\}$) such that for each pair of different colors $i, j \in [k]$ there

*Research supported by CONACyT-México under Projects 57371, 166306, PAPIIT-México under Project IN104915.

exists an edge $xy \in E(G)$ where $x \in \varsigma^{-1}(i)$ and $y \in \varsigma^{-1}(j)$. The *pseudoachromatic number* $\psi(G)$ of G is the largest k for which there exists a complete k-coloring of G [11]. Some interesting results on the pseudoachromatic number and the closely related notion of the achromatic number (the maximum on proper and complete colorings) can be found in [6, 8, 13].

The *connected-pseudoachromatic number* $\psi_c(G)$ of a connected graph G is the largest k for which there exists a connected and complete k-coloring of G, i.e., a complete k-coloring in which each color class induces a connected subgraph. The connected pseudoachromatic number $\psi_c(G)$ of a graph G with components G_1,G_2,...,G_t is the largest number between $\psi_c(G_1)$,$\psi_c(G_2)$,...,$\psi_c(G_t)$. Clearly,

$$\psi_c(G) \leq \psi(G).$$

The previous definition is equivalent to say that $\psi_c(G)$ is the size of the largest complete graph minor of G. This value is also called the *Hadwiger number* of G. In 1958, Hadwiger conjectured (see [12]) that for any G graph $\chi(G) \leq \psi_c(G)$ where $\chi(G)$ denotes, as usual, the chromatic number of G and therefore. Aside from their own importance the Hadwiger Conjecture is another motivation to study the previous parameters since the pseudoachromatic number bounding the Hadwiger number.

The connected-pseudoachromatic and the pseudoachromatic numbers of the line graph $L(G)$ of a graph G are also known as the *connected-pseudoachromatic index* and the *pseudoachromatic index* of G denoted as $\psi_c'(G)$ and $\psi'(G)$ respectively. In this paper, we study these two parameters for the complete graph K_n. Note that any connected and complete k-coloring of $L(K_n)$ is an edge coloring of K_n in which each edge color class induces a connected subgraph and each pair of edge color classes share at least one vertex, therefore, we will make use of this point of view.

It is important to note that the colorations that induce the lower bound are given using the structure of projectives planes, it is a common technique used before to obtain results of these two parameters and also for the achromatic index of the complete graph K_n, this could be confirmed in many of the following references related to these topics [1, 3, 8, 4, 9, 15]

Before continuing, we give some known results on $\psi_c'(K_n)$ and $\psi'(K_n)$ related to our contributions. On the case of exact values of $\psi_c'(K_n)$ only the following small values of Table 1 are known. Table 1 appears in [1], which also proves that $\psi_c'(K_{5a+b+1})$ is at least $9a + b$.

n	2	3	4	5	6	7
$\psi_c'(K_n)$	1	3	4	6	7	10

Table 1: Exact values for $\psi_c'(K_n)$, $2 \leq n \leq 7$.

However, in Table 2 (which appears also in [3]) the first pseudoachromatic index values of the complete graph are enlisted.

n	2	3	4	5	6	7	8	9	10	11	12	13
$\psi'(K_n)$	1	3	4	7	8	11	14	18	22	27	32	39

Table 2: Exact values for $\psi'(K_n)$, $2 \leq n \leq 13$.

The pseudoachromatic index for a few classes of complete graphs have been found so far, namely, Bouchet proved in [9] that for $n = q^2 + q + 1$ and q an odd integer $\alpha'(n) = qn$, if and only if the projective plane of order q exists, in consequence, it is known $\psi'(K_{q^2+q+1})$ for any odd prime power q (see [4, 9]). On the other hand, when q is a power of 2 the pseudoachromatic index is bounding by $\psi'(K_{q^2+q+1}) \geq q^3 + q$ (see [4]) and attains exact values for $K_{q^2+2q+1-a}$ when $a \in \{-1, 0, \ldots, \left\lceil \frac{1+\sqrt{4q+9}}{2} \right\rceil - 1\}$ (see [2, 3, 4]). Finally, it is proved that $\psi'(K_n)$ grows asymptotically like $n^{3/2}$ (see [15]).

This paper is organized as follows. In the first section, we give some definitions and lemmas related to projective planes and colorings which will be used in the second and third sections in order to give lower bounds for the pseudoachromatic connected and pseudoachromatic indices of complete graphs. In Section 2, we prove that, for any integer $n \geq 8$, $\psi'_c(K_n)$ has order $n^{3/2}$. The proof is divided in the following results:

Theorem 1. *If $n \geq 8$ then*

$$\psi'_c(K_n) \leq \lfloor \max \{\min\{f_n(x), g_n(x) : x \in \mathbb{N}\}\} \rfloor$$

where $f_n(x) = n(n-1)/2x$ and $g_n(x) = (x+1)(n-x-1/2)$.

Theorem 2. *If $n \geq 8$ then*

$$\psi'_c(K_n) \leq \frac{1}{\sqrt{2}} n^{\frac{3}{2}} + \Theta(n).$$

Theorem 3. *If q is a prime power and $n = q^2 + q + 1$ then*

$$\left\lceil \frac{q}{2} \right\rceil n \leq \psi'_c(K_n).$$

Theorem 4. *If $n \geq 8$ then*

$$\frac{1}{2} n^{\frac{3}{2}} + \Theta(n) \leq \psi'_c(K_n).$$

Note that comparing the results of Theorems 1 and 2 with the upper bounds for the pseudoachromatic number given in [2, 3, 4, 15] we have that $\psi'_c(K_n) < \psi'(K_n)$. On the other hand Theorems 3 and 4 improve the linear lower bound that appears in [1] and verifies the Hadwiger Conjecture. Finally, Theorems 2 and 4 show that $\psi'_c(K_n)$ grows asymptotically like $\Theta(n^{3/2})$.

In the third section we study the pseudoachromatic index $\psi'(K_n)$ of the complete graph K_n for $n = q^2 + q + 1$ and q a power of 2. Specifically, we improve the lower bound that appears in [4] proving the following theorem:

Theorem 5. *If q is a power of 2 and $n = q^2 + q + 1$ then*

$$q^3 + 2q - 3 \leq \psi'(K_n).$$

2 Preliminaries

A *projective plane* consists of a set of n points, a set of lines, and an incidence relation between points and lines having the following properties:

1. Given any two distinct points there is exactly one line incident with both.
2. Given any two distinct lines there is exactly one point incident with both.
3. There are four points, such that no line is incident with more than two of them.

Such plane has $n = q^2 + q + 1$ points (for some number q) and n lines, each line contains $q + 1$ points and each point belongs to $q + 1$ lines. The number q is called the *order* of the projective plane. A projective plane of order q is called Π_q. If q is a prime power there exists Π_q, which is called the *algebraic projective plane* since it arises from finite fields (see [14]).

Let $\mathbb{P}$ be the set of points of Π_q and let $\mathbb{L} = \{l_1, \ldots, l_n\}$ the set of lines of Π_q. Now identify the points of Π_q with the set of vertices of the complete graph K_n. In a natural way the set of points of each line of Π_q induces in K_n a subgraph isomorphic to K_{q+1}. For each line $l_i \in \mathbb{L}$, let $l_i = (V(l_i), E(l_i))$ be the subgraph of K_n induced by the set of $q + 1$ points of l_i. By the properties of the projective plane, for every pair $\{i, j\} \subseteq \{1, \ldots, n\}$, $|V(l_i) \cap V(l_j)| = 1$ and $\{E(l_1), \ldots, E(l_n)\}$ is a partition of $E(K_n)$. In this manner when we say that a graph G isomorphic to K_n is a *representation of the projective plane* Π_q we will understand that $V(G)$ is identified with the points of Π_q and that there is a family of subgraphs (lines) $\{l_1, \ldots, l_n\}$ of G, such that for each line l_i of Π_q, l_i is the subgraph induced by the set of points of l_i. This is a standard method of correspondence.

Given two graphs G and H, the *join* $G \oplus H$, is defined as the graph with vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H) \cup (V(G) \times V(H))$ (to abbreviate we write the set of edges $V(G) \times V(H)$ as $V(G)V(H)$-edges). Given $S \subseteq V(G)$, $G \setminus S$ is the subgraph of G induced by $V(G) \setminus S$.

Given an edge-coloring $\varsigma \colon E(K_n) \to [k]$ we will say that a vertex $x \in V(K_n)$ is an *owner* of a set of colors $\mathcal{C} \subseteq [k]$ whenever for every $i \in \mathcal{C}$ there is $y \in V(K_m)$, such that $\varsigma(xy) = i$; and given a subgraph G of K_n we will say that G is an *owner* of a set of colors $\mathcal{C} \subseteq [k]$, if each vertex of G is an owner of $\mathcal{C}$. By this ς is a complete edge-coloring, if for every pair of colors in $[k]$ there is a vertex in K_n, which is an owner of both colors.

Lemma 1. *Let $n = q^2 + q + 1$ with q a natural number, such that Π_q exists. Let K_n be a representation of Π_q and let $\varsigma \colon E(K_n) \to [k]$ an edge-coloring of K_n. Suppose that each line l_i of K_n is an owner of a set of colors $\mathcal{C}_i \subseteq [k]$. Then for every pair of colors $\{c_1, c_2\} \subseteq \bigcup_{i=1}^{n} \mathcal{C}_i$ there is $x \in V(G)$, which is an owner of c_1 and c_2.*

Proof. Let $\{c_1, c_2\} \subseteq \bigcup_{i=1}^{n} \mathcal{C}_i$. If there is an $i \in \{1, \ldots, n\}$, such that $\{c_1, c_2\} \subseteq \mathcal{C}_i$ then since l_i is the owner of $\mathcal{C}_i$ it follows that each $x \in V(l_i)$ is an owner of c_1 and c_2. If $c_1 \in \mathcal{C}_i$ and $c_2 \in \mathcal{C}_j$ with $i \neq j$ there is $x \in V(G)$, such that $x = V(l_i) \cap V(l_j)$ and then since l_i and l_j are owners of $\mathcal{C}_i$ and $\mathcal{C}_j$ respectively, x is an owner of c_1 and c_2. ∎

It is well known that any complete graph of even order $q + 1$ admits a 1-factorization $\mathcal{M}$ by q perfect matchings and that any complete graph of odd order $q + 1$ admits a 2-factorization $\mathcal{H}$ by $q/2$ Hamiltonian cycles (see [7]).

Using the above, we define different edge-colorations for some special graphs. The first two will be used to give the results stated in Theorem 3 whereas the last four will be necessary to prove the results of Theorem 5.

Definition 1. Let $q+1$ be an odd integer. An edge-coloring $\varsigma\colon E(K_{q+1}) \to [q/2]$ will be said to be of *Type* $\mathcal{H}$, if for every $i \in [q/2]$ the set $\{xy \in E(K_{q+1})\colon \varsigma(xy) = i\}$ is a Hamiltonian cycle of K_{q+1}.

Definition 2. Let $q+1$ be an even integer. An edge-coloring $\varsigma\colon E(K_{q+1}) \to [(q+1)/2]$ will be said to be of *Type* $\mathcal{P}$, if we obtain ς in the following way: Let G be the graph (isomorphic to K_{q+2}) obtained by adding a new vertex x_0 and all the $x_0V(K_{q+1})$-edge. Let ς_1 be an edge-coloring of Type $\mathcal{H}$ of G and for every $e \in E(K_{q+1})$ let $\varsigma(e) := \varsigma_1(e)$. Note that for every $i \in [(q+1)/2]$ the set $\{xy \in E(K_{q+1})\colon \varsigma(xy) = i\}$ is a Hamiltonian path of K_{q+1}.

Definition 3. Let $q+1$ be an odd integer. An edge-coloring $\varsigma\colon E(K_{q+2}) \to [q+1]$ is called of *Type* $\mathcal{M}$, if for every $i \in [q+1]$ the set of edges $\varsigma^{-1}(i)$ is a perfect matching of K_{q+2}.

Definition 4. Let $q+1$ be an odd integer. An edge-coloring $\varsigma\colon E(K_{q+1} - xy) \to [q-1]$ will be said to be of *Type* $\mathcal{C}$, if we obtain ς in the following way: Let G be the graph (isomorphic to K_q) obtained by deleting the vertex x and all the $xV(K_q)$-edges. Let ς_1 be an edge-coloring of Type $\mathcal{M}$ of G and, for every $e \in E(K_{q+1} - xy)$, let $\varsigma(e) := \varsigma_1(e)$, if $e \in E(G)$ and $\varsigma(xw) := \varsigma_1(yw)$ for all $w \in V(G) - y$. The edge xy is called the *special edge* of K_{q+1} (see Figura 1).



Figure 1: An edge-coloring of Type $\mathcal{C}$ for $K_5 - xy$. The special edge xy is dashed.

Definition 5. Let $q+1$ be an odd integer. An edge-coloring $\varsigma : E(K_{q+1} - M) \to [q]$ will be called of *Type 1*, if we obtain ς in the following way: Let G be the graph (isomorphic to K_{q+2}) obtained by adding a new vertex x and all the $xV(K_{q+1})$-edges. Let $\varsigma_1\colon E(G) \to [q+1]$ an edge-coloring of Type $\mathcal{M}$ of G. For every $e \in E(K_{q+1})$, $\varsigma(e) := \varsigma_1(e)$, and M is a maximum matching of K_{q+1}, such that $E(M) \subseteq \varsigma_1^{-1}(q+1)$. Note that there exist a vertex y of $K_{q+1} - M$, such that $M \cup \{xy\} = \varsigma_1^{-1}(q+1)$ is a perfect matching of G (see Figura 2).

By the construction of edge-coloring of Type 1, the vertex y of $K_{q+1} - M$ is owner of its q colors, but the remaining vertices are owner of its q colors except for one. Moreover, for all $x \in K_{q+1} - M$ and $x \neq y$ we will call the *missing color of* x the only color for which x is not owner. The following edge-coloring makes K_{q+1} owner of the q colors.

Definition 6. Let $q+1$ be an odd integer, K_{q+1} the complete graph of order $q+1$ with the set of vertices $\{y, x_1, x_1', \ldots, x_{q/2}, x_{q/2}'\}$, $M = \{x_1x_1', \ldots, x_{q/2}x_{q/2}\}$ a maximum matching of K_{q+1}, and let G be the graph obtained by adding a new vertex x and the set of edges $\{xx_1, \ldots, xx_{q/2}\}$. An edge-coloring $\varsigma : E(G) \to [q]$ will be called of *Type 2*, if we obtain ς in the following way: Let $\varsigma_1\colon E(K_{q+1} - M) \to [q]$ be an edge-coloring of Type 1 of the subgraph $K_{q+1} - M$ of G. For every $e \in E(K_{q+1}) - M$, $\varsigma(e) := \varsigma_1(e)$. If c_i is the missing color of x_i and c_i' is the missing color of x_i' (for all $i \in [q/2]$) let $\varsigma(xx_i) = c_i$ and $\varsigma(x_ix_i') = c_i'$. The vertex x is called the *special vertex* of G (see left Figure 2).



Figure 2: Left: An edge-coloring of Type 1 for $K_5 - M$, the maximum matching M is dashed. Right: An edge-coloring of Type 2 for G.

3 Connected-pseudoachromatic index of K_n

We start the section proving Theorems 1 and 2, these results provide the upper bound of the connected-pseudoachromatic index of K_n. The proofs are analytic and their main idea is to compare two functions one of which determines how many chromatic classes of size x there are at most and the other one determines how many chromatic classes can be incident in one chromatic class of size x.

Proof of Theorem 1. Let $n \geq 8$ and $\varsigma\colon E(K_n) \to [k]$ a connected and complete k-edge-coloring of K_n with $k = \psi'_c(K_n)$. Let $x = min\{\left|\varsigma^{-1}(i)\right| : i \in [k]\}$, that is, let x be the cardinality of the smallest chromatic class of ς and suppose that $x = \left|\varsigma^{-1}(k)\right|$. Since ς defines a partition of $E(K_n)$ it follows that $k \leq f_n(x) := n(n-1)/2x$. On one hand, since ς is connected we can suppose that $\left\langle \varsigma^{-1}(k)\right\rangle$ is a tree. On the other hand, since ς is complete, for every $j \in [k-1]$ there is a vertex $x_j \in \varsigma^{-1}(j)$, such that x_j is adjacent with some edge in $\varsigma^{-1}(k)$ and since the number of trees in $\left\langle E(K_m) \setminus \varsigma^{-1}(k)\right\rangle$, which are adjacent to some edge in $\varsigma^{-1}(k)$ is at most

$$g_n(x) - 1 := (x+1)(n-(x+1)) + \frac{\binom{x+1}{2} - x}{x},$$

it follows that $k \leq g_n(x) = (x+1)(n-x-1/2)$. Therefore,

$$\psi'_c(K_n) \leq \lfloor \max\{\min\{f_n(x), g_n(x) : x \in \mathbb{N}\}\}\rfloor .$$

■

In order to prove Theorem 2 we need the following technical lemma:

Lemma 2. *Let $x_0, x_1 \in \mathbb{R}^+$, such that $f_n(x_0) = g_n(x_0)$ and $f_n(x_1) = g_n(x_1)$ and suppose that $x_0 < x_1$, then*

$$g_n(x_0) = f_n(x_0) = \max\left\{\min\{f_n(x), g_n(x) : x \in \mathbb{R}^+\}\right\}$$

Furthermore, $x_0 = \sqrt{n/2 + 1/16} - 1/4$.

Proof. As f_n is a hyperbola and g_n is a parabola (see Figura 3) it is not hard to see that $f_n(x) \leq g_n(x)$ for $x_0 \leq x \leq x_1$ and $g_n(x) < f_n(x)$ in any other case when $x \in \mathbb{R}^+$. Since $f_n(x_0) > f_n(x_1)$ it follows that $g_n(x_0) = f_n(x_0) = \max\{\min\{f_n(x), g_n(x) : x \in \mathbb{R}^+\}\}$.

Finally, the equation $f_n(x_0) = g_n(x_0)$ is reduced to $n^2 - (2x_0^2 + 2x_0 + 1)n + x_0(x_0+1)(2x_0+1) = 0$. Taking the positive solution for n we get $n = 2x_0^2 + x_0$. It is not difficult to prove that $x_0 = \sqrt{n/2 + 1/16} - 1/4$ provides the positive solution. ■



Figure 3: The functions g_n and f_n for some fixed value n.

Proof of Theorem 2. Since $g_n(x_0) = nx_0 - x_0^2 - 3x_0/2 + n - 1/2$ and $x_0 = \sqrt{n/2 + 1/16} - 1/4$ we obtain

$$g_n(x_0) = (n-1)(\sqrt{n/2 + 1/16} + 1/4),$$

therefore $g_n(x_0) \leq \frac{1}{\sqrt{2}} n^{\frac{3}{2}} + \Theta(n)$. By Lemma 2 and Theorem 1 the result follows. ∎

Now we will proof Theorem 3 to give a lower bound for the complete graphs with some specific order related to the projective planes.

Proof of Theorem 3. Suppose q is the order of a projective plane Π_q. Then Π_q has $n = q^2 + q + 1$ points and n lines. Let K_n be a representation of Π_q. Now we color K_n in the following way: if q is odd, each line K^i_{q+1} has a coloring $\varsigma_i \colon E(K^i_{q+1}) \to \{1 + (i-1)q/2, \ldots, q/2 + (i-1)q/2\}$ of Type $\mathcal{H}$ (for $i \in [n]$). Each line K^i_{q+1} may thus be divided into $q/2$ color classes, each one of size $q+1$ and it is owner of its colors. By Lemma 1 the $qn/2$-edge-coloring is complete and by construction is connected. Similarly, if q is even, K_n has a connected and complete $(q+1)n/2$-edge-coloring using edge-coloring of Type $\mathcal{P}$ in its lines. ∎

Lemma 3. *For any connected graph G, if H is a connected subgraph of G, then $\psi_c(G) \geq \psi_c(H)$.*

Proof. Given a connected and complete coloring ς of H with $\psi_c(H)$ colors, we extend this to a coloring of G by coloring the set of vertices of a component G' of $G \setminus H$ with some color that appears on a vertex of H adyacent to a some vertex of G'. ∎

Proof of Theorem 4. The proof depends on a strengthened version of Bertrand's "Postulate" which follows from the Prime Number Theorem (see [5, 10]): For any $\epsilon > 0$, there exists an N_ϵ such that for any real $x \geq N_\epsilon$, there is a prime q between x and $(1+\epsilon)x$. Now, let $\epsilon > 0$ given, and suppose $n > (N_\epsilon + 1)^2(1+\epsilon)^2$. Let $x = \sqrt{n}/(1+\epsilon) - 1$, so $x \geq N_\epsilon$. We may then select a prime q with $x \leq q \leq (1+\epsilon)x$. Note that $q^2 + q + 1 \leq (x+1)^2(1+\epsilon)^2 = n$. Since projective planes of all prime orders exist, it follows from Theorem 3 and Lemma 3 that

$$\psi'_c(K_n) \geq \psi'_c(K_{q^2+q+1}) \geq \frac{q}{2}(q^2 + q + 1) > \frac{q^3}{2} \geq \frac{x^3}{2} = \frac{(\sqrt{n} - 1 - \epsilon)^3}{2(1+\epsilon)^3}.$$

Since ϵ was arbitrarily small, the result follows. ∎

4 Pseudoachromatic index of some complete graphs related to projective planes of even order

As in the previous proofs, for the proof of the last theorem, it is enough to exhibit a complete (q^3+2q-3)-coloring of K_{q^2+q+1} for q a power of two, since the projective plane of order q always exists.

Proof of Theorem 5. To prove this theorem we will exhibit a complete edge-coloring of K_n for $n = q^2+q+1$, with $k := q^3+2q-3$ colors. Let C be a set of k colors and let $\{C_1, C_2, \ldots, C_n\}$ be a partition of C in the following way: C_i is a set of $q-1$ colors, for $1 \leq i \leq q^2-q+3$; and C_i is a set of q colors, for $q^2-q+4 \leq i \leq n$.

Let G be a graph isomorphic to K_n, let G be a representation of Π_q and let $L = \{l_1, \ldots, l_n\}$ the set of lines of G.

Let $V(l_n) = \{p_1, \ldots, p_{q+1}\}$. For $i \in [q+1]$, let $L_i = \{l_{q(i-1)+j} : p_i \in V(l_{q(i-1)+j}), \forall j \in [q]\} \subseteq L \backslash \{l_n\}$.

We start to coloring the edges of K_n with $\varsigma\colon E(K_n) \to [k]$ in the following way:

i) For $i \in \{0,1,2\}$, let $\varsigma_{n-i} : E(l_{n-i} - e_{n-i}) \to C_{q^2-q+3-i}$ an edge-coloring of Type $\mathcal{C}$ with the special edge e_{n-i} (any edge of l_{n-i}).

ii) For $i \in \{1, \ldots, q-1\}$, let $\varsigma_{q(i-1)+j} : E(l_{q(i-1)+j} - e_{q(i-1)+j}) \to C_{q(i-1)+j}$ an edge-coloring of Type $\mathcal{C}$ with the special edge $e_{q(i-1)+j} := p_i x_{q(i-1)+j}$ where $x_{q(i-1)+j}$ depends of the values of i and j:

$$x_{q(i-1)+j} := \begin{cases} l_{q(i-1)+j} \cap l_{q(q-1)+2i-1} & \text{if } i,j \in \{1, \ldots, q/2\} \\ l_{q(i-1)+j} \cap l_{q(q-1)+2i} & \text{if } i \in \{1, \ldots, q/2\} \text{ and } j \in \{q/2+1, \ldots, q\} \\ l_{q(i-1)+j} \cap l_{q^2+2i-1} & \text{if } i \in \{q/2+1, \ldots, q-1\} \text{ and } j \in \{1, \ldots, q/2\} \\ l_{q(i-1)+j} \cap l_{q^2+2i} & \text{if } i \in \{q/2+1, \ldots, q-1\} \text{ and } j \in \{q/2+1, \ldots, q\}. \end{cases}$$

Figure 4 depicts the first and the second cases for $i = 1$.

iii) Let $G_j := (V(G_j), E(G_j))$ be a graph, then we have the following two subcases:

- If $V(G_j) = V(l_{q(q-1)+j}) \cup \{p_{\lceil j/2 \rceil}\}$ and $E(G_j) = E(l_{q(q-1)+j}) \cup \{e_{(j-1)q/2+1}, \ldots, e_{(j-1)q/2+q/2}\}$ with $j \in \{1, \ldots, q\}$; let $\varsigma_{q(q-1)+j} : E(G_j) \to C_{q^2-q+3+j}$ be an edge-coloring of Type 2 with the special vertex $p_{\lceil j/2 \rceil}$.



Figure 4: A drawing of some lines l_j and some vertices x_j for $i = 1$.

- And, if $V(G_j) = V(l_{q^2+j}) \cup \{p_{q/2+\lceil j/2 \rceil}\}$ and $E(G_j) = E(l_{q^2+j}) \cup \{e_{q^2/2+(j-1)q/2+1}, \ldots, e_{q^2/2+(j-1)q/2+q/2}\}$ with $j \in \{1, \ldots, q-2\}$; let $\varsigma_{q^2+j} : E(G_j) \to C_{q^2+3+j}$ be also an edge-coloring of Type 2 with the special vertex $p_{q/2+\lceil j/2 \rceil}$.

iv) We have already assigned a color to each edge of $E(K_n) \backslash \{e_n, e_{n-1}, e_{n-2}\}$. Let coloring the set of 3 edges $\{e_n, e_{n-1}, e_{n-2}\}$ with the color 1.

By the construction and Lemma 1 it follows that the resultant edge-coloring $\varsigma := \bigcup_{i=1}^{n} \varsigma_i$ of K_n is a complete edge-coloring, since every line l_i in K_n is an owner of its set of colors. ■

References

[1] L. Abrams and Y. Berman. Connected pseudoachromatic index of complete graphs. *Australas. J. Combin.*, 60:314–324, 2014.

[2] G. Araujo-Pardo, J. J. Montellano-Ballesteros, C. Rubio-Montiel, and R. Strausz. On the pseudoachromatic index of the complete graph III. *In review.*

[3] G. Araujo-Pardo, J. J. Montellano-Ballesteros, C. Rubio-Montiel, and R. Strausz. On the pseudoachromatic index of the complete graph II. *Bol. Soc. Mat. Mex. (3)*, 20(1):17–28, 2014.

[4] G. Araujo-Pardo, J. J. Montellano-Ballesteros, and R. Strausz. On the pseudoachromatic index of the complete graph. *J. Graph Theory*, 66(2):89–97, 2011.

[5] B. Bollobás. *Extremal graph theory*, volume 11 of *London Mathematical Society Monographs*. Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], London-New York, 1978.

[6] B. Bollobás, B. Reed, and A. Thomason. An extremal function for the achromatic number. In *Graph structure theory (Seattle, WA, 1991)*, volume 147 of *Contemp. Math.*, pages 161–165. Amer. Math. Soc., Providence, RI, 1993.

[7] J. A. Bondy and U. S. R. Murty. *Graph theory with applications*. American Elsevier Publishing Co., Inc., New York, 1976.

[8] J. Bosák and J. Nešetřil. Complete and pseudocomplete colourings of a graph. *Math. Slovaca*, 26(3):171–184, 1976.

[9] A. Bouchet. Indice achromatique des graphes multiparti complets et réguliers. *Cahiers Centre Études Rech. Opér.*, 20(3-4):331–340, 1978.

[10] A. A. Gioia. *The theory of numbers. An introduction*. Markham Publishing Co., Chicago, Ill., 1970.

[11] R. P. Gupta. Bounds on the chromatic and achromatic numbers of complementary graphs. In *Recent Progress in Combinatorics (Proc. Third Waterloo Conf. on Comb., 1968)*, pages 229–235. Academic Press, New York, 1969.

[12] H. Hadwiger. Ungelöste probleme 26. *Elem. Math.*, 13:128–129, 1958.

[13] F. Harary, S. Hedetniemi, and G. Prins. An interpolation theorem for graphical homomorphisms. *Portugal. Math.*, 26:453–462, 1967.

[14] J. W. P. Hirschfeld. *Projective geometries over finite fields*. The Clarendon Press, Oxford University Press, New York, 1979. Oxford Mathematical Monographs.

[15] R. E. Jamison. On the edge achromatic numbers of complete graphs. *Discrete Math.*, 74(1-2):99–115, 1989.